

Cite this: *CrystEngComm*, 2014, 16, 5364

# Influence of the anion nature on styryl dye crystal packing and feasibility of the direct and back [2 + 2] photocycloaddition reactions without single crystal degradation†

 Lyudmila G. Kuz'mina,<sup>\*a</sup> Artem I. Vedernikov,<sup>b</sup> Andrei V. Churakov,<sup>a</sup> Elmira Kh. Lermontova,<sup>a</sup> Judith A. K. Howard,<sup>c</sup> Michael V. Alfimov<sup>b</sup> and Sergey P. Gromov<sup>\*b</sup>

A family of styryl dyes of the 4-pyridine series,  $\text{Et-Py}^+-\text{CH}=\text{CH}-\text{C}_6\text{H}_3(\text{OMe})_2 \text{X}^-$ , with different anions  $\text{X}^-$ , was synthesized. Their ability to undergo [2 + 2] photocycloaddition (PCA) in the polycrystalline state with the formation of 1,2,3,4-tetrasubstituted *rc*tt cyclobutanes under irradiation with visible light was investigated by  $^1\text{H}$  NMR spectroscopy. Crystal packing of the dyes and feasibility for the PCA reaction in single crystals were investigated by X-ray structural analysis. Small inorganic anions, such as  $\text{ClO}_4^-$ ,  $\text{BF}_4^-$ ,  $\text{PF}_6^-$  that do not form secondary interactions with the organic cation, aid the formation of cation stacking motifs of the "head-to-tail" type. In these stacks, any adjacent cations are arranged in an antiparallel fashion and approach one another at distances shorter than 4.2 Å, which corresponds to the Schmidt's rule for feasibility of PCA reaction in the solid state. As a result of PCA, *rc*tt isomers of cyclobutane derivatives are formed. Success of the PCA reaction without single crystal degradation was only found for the dye containing  $\text{BF}_4^-$  anion. Upon irradiation with UV light of the cyclobutane product formed in the crystal, back phototransformation without single crystal degradation was observed. A number of weak interactions  $\text{I}^-\cdots\text{H}-\text{C}$  in the dye with  $\text{I}^-$  anion prevent significant atomic displacements that accompany the PCA transformation from being achieved. Occurrence of the bulky tosylate anion forming a set of hydrogen bonds with solvate acetonitrile and water molecules aids the formation of "head-to-head" type stacks, with a large separation between the adjacent ethylene bonds. In the dye containing bulky  $\text{BPh}_4^-$  anion, the main structure-forming role belongs to anions, which suppress the cation trend to form a stacking packing motif. The dye with picrate anion forms two types of crystals. Crystal packing of one of them excludes any PCA transformation, whereas, the other one permits it, but with single crystal degradation.

 Received 4th February 2014,  
Accepted 26th March 2014

DOI: 10.1039/c4ce00266k

[www.rsc.org/crystengcomm](http://www.rsc.org/crystengcomm)

## Introduction

The solid phase reaction of [2 + 2] photocycloaddition (PCA) of unsaturated compounds resulting in cyclobutane derivatives has created great interest since the 1960's. On the one

hand this reaction is the best way for obtaining substituted cyclobutanes<sup>1–20</sup> and on the other hand, it may be used in systems for optical data recording due to the considerable change in physico-chemical properties of the compounds.<sup>21,22</sup>

The PCA reaction in these compounds is accomplished upon their irradiation with visible or near UV-range light in both solid and liquid phases. The PCA reaction without single crystal degradation presents a special case.<sup>3–5,7,9,12–14,20,23–29</sup> However, more frequently the reaction leads to crystal degradation to amorphous glass-like state or powder. The reasons why a single crystal retains or decomposes in the course of PCA are not always evident, although in individual cases they become evident from the analysis of X-ray structural data.<sup>23–29</sup>


For the PCA reaction to occur, a particular spatial pre-organization of two starting unsaturated molecules is required (Scheme 1). The molecules should be arranged in parallel

<sup>a</sup> Institute of General and Inorganic Chemistry, Russian Academy of Sciences, Leninskii prosp. 31, Moscow 119991, Russian Federation. E-mail: kuzmina@igic.ras.ru; Fax: +7 (495) 954 1279

<sup>b</sup> Photochemistry Centre, Russian Academy of Sciences, Novatorov str. 7A-1, Moscow 119421, Russian Federation. E-mail: spgromov@mail.ru; Fax: +7 (495) 936 1255

<sup>c</sup> Department of Chemistry, University of Durham, Science Laboratories, South Road, Durham DH1 3LE, England, UK. E-mail: j.a.k.howard@durham.ac.uk

† CCDC 969548–969552 and 859813–859815. For crystallographic data in CIF or other electronic format see DOI: 10.1039/c4ce00266k




Scheme 1 View of preorganization of two ethylene units for PCA reaction.

planes in such a way that their ethylene groups are located one above another and are oriented parallel (or antiparallel) to each other with the distance  $d$  between the carbon atoms varying in the range 3.4–4.2 Å (Schmidt's rule).<sup>1</sup>

This preorganization can be achieved as a result of a molecular design. In styryl dyes of general formula  $R\text{-Het}^+-\text{CH}=\text{CH-Ar}^-$  or their precursors – neutral styrylheterocycles  $\text{Het-CH}=\text{CH-Ar}$  – the actual geometry of the initial preorganized dimer can be controlled by involving different functional fragments in ethylene-containing systems. For instance, one of these fragments may be a benzocrown-ether group forming sandwich dimers in the presence of big metal cations in solutions,<sup>30</sup> whereas an addition of a *N*-sulfonatoalkyl or *N*-ammonioalkyl group in the opposite position of this molecular system aids the formation of another geometry of the preorganized dimers in the solid state and in solution (Scheme 2).<sup>31–34</sup>

The creation of the other type of sandwich dimers *via* binding to a metal cation or hydrogen bonds forming between the components<sup>3,5–7,9,10,12–16,18–20,35</sup> or due to the assistance of macroheterocyclic compounds (crown ethers, cucurbit[8]uril,  $\gamma$ -cyclodextrin, calix[*n*]arenes, *etc.*)<sup>8,28,36–44</sup> is also described for unsaturated compounds.




Scheme 2 The structure of dimers with different types of preorganization for PCA.

In this work, we used another approach for creating sandwich dimers, taking into account that the crystal lattice may perform a topochemical control over the PCA reaction, namely, the requirements for PCA preorganized dimers are enhanced by their crystal environment.<sup>23–27,29,45</sup> Then the investigation of PCA reactions is based on crystal engineering.<sup>46</sup> The investigation involves analyzing the crystal packing for the group of compounds related to those chosen for study and searching for ways to construct the optimal crystal packing.

Previously, we have investigated the influence of the cation and the occurrence of additional building blocks in the crystal packing of the related styryl dyes on the formation of a particular crystal packing favourable for the PCA reaction without single crystal degradation.<sup>23–27,29,45</sup> In the last case, relatively small aromatic solvent molecules were used as additional building blocks.<sup>27</sup> The aim of this work is to explore possibilities for controlling crystal packing motifs by anion variation. Dyes of the 4-pyridine series **1a–g** chosen as objects of this study are shown in Scheme 3.

These anions have the same charge but different shapes, spatial behaviors and abilities to form weak interactions with surrounding functional groups in the crystal. These are the spherical shape anion ( $\text{I}^-$ ), pseudo-spherical shape anions revealing a fluxional behavior resulting in a rotational disorder in crystals ( $\text{ClO}_4^-$ ,  $\text{BF}_4^-$ ,  $\text{PF}_6^-$ ), the non-symmetrical anion having conformational freedoms ( $\text{TsO}^-$ ), the big planar anion prone to form a charge-transfer complex (CTC) with aromatic species ( $\text{Pic}^-$ ), and the bulky nonplanar anion  $\text{BPh}_4^-$ . The two last anions are capable of influencing the crystal packing formation significantly.

Here we describe the synthesis and X-ray structure determination of dyes **1c–g** and analyze the crystal packing for series **1c–g**, as well as the previously investigated dyes **1a** and **1b**.<sup>25,26</sup> After the X-ray study of the compounds was carried out, their single crystals were subjected to irradiation with visible light. For the crystals that are saved unchanged, the second X-ray structural study was performed. For compound **1d** that is found to transform into a cyclobutane derivative without single crystal degradation, we attempted to record a back (retro) photochemical reaction without single crystal degradation by irradiating the cyclobutane product of the direct photochemical reaction with UV light. For comparison, we also performed a visible light irradiation of thin polycrystalline films of dyes **1a–g** and analyzed the composition of products and the degree of photoconversion using  $^1\text{H}$  NMR.



Scheme 3 The structure formula of dyes **1**:  $\text{X}^- = \text{ClO}_4^-$  (a),  $\text{I}^-$  (b),  $\text{TsO}^-$  (c),  $\text{BF}_4^-$  (d),  $\text{PF}_6^-$  (e),  $\text{Pic}^-$  (picrate anion) (f), and  $\text{BPh}_4^-$  (g).


## Results and discussion

### Synthesis and photochemical investigation in polycrystalline films

The synthesis of dyes **1a** and **b** has been described earlier.<sup>25</sup> The synthesis of dye **1c** containing the tosylate anion was performed by quaternization of neutral dimethoxystyrylpyridine with ethyl 4-methylbenzenesulfonate. Dyes **1d–g** were obtained by anion exchange in **1c** on treatment with the appropriate salt (NaBF<sub>4</sub>, NH<sub>4</sub>PF<sub>6</sub>, NaPic, NaBPh<sub>4</sub>) in methanol (Scheme 4). All new compounds were characterized by <sup>1</sup>H NMR spectroscopy data and elemental analysis.

Yellow dyes **1a–g** in the form of polycrystalline films on glass substrates were irradiated with visible light under comparable conditions. The compositions of the samples thus obtained were analyzed by <sup>1</sup>H NMR spectroscopy. The results are given in Table 1.

Dye **1g** proved to be insensitive to light. By contrast, the irradiation of the other compounds resulted in the formation



Scheme 5 Synthesis of cyclobutane derivatives *rctt*-2a–f.

of centrosymmetric *rctt* isomers of cyclobutane derivatives **2a–f** (Scheme 5). The <sup>1</sup>H NMR spectra of **2a–f** show characteristic<sup>23–25,28,29</sup> signals of cyclobutane protons at  $\delta$  4.8–4.9: two symmetric doublets of doublets and the AA'BB'-type spin system. This is indicative of the existence of *syn*-‘‘head-to-tail’’ stacked dimers in the solid state, which should precede the formation of cyclobutanes with this stereochemistry. The rates of the PCA reactions are substantially different. The phototransformation proceeds most rapidly in dyes **1a**, **c–e** (the degree of conversion is 87–100% for 10 h), whereas the complete conversion of picrate **1f** was not achieved even within >240 h. The phototransformation of iodide **1b** is also found to be much slower.<sup>25</sup>

In order to check a possibility for the solid state *retro*-PCA reaction in the compounds obtained, we performed the following experiment. At first, a polycrystalline film of dye **1d** was irradiated with visible light ( $\lambda > 390$  nm) until full transformation to cyclobutane derivative *rctt*-**2d** (monitored by <sup>1</sup>H NMR). Then this film was subjected to short-wave irradiation ( $\lambda$  in the range of 280–340 nm). Fragments of the <sup>1</sup>H NMR spectra of dye **1d** and photolyzates of its film are shown in Fig. 1.


It is observed that on irradiation of the initial compound, its proton signals in the spectrum have disappeared completely, whereas, the subsequent UV irradiation results in their reappearance; the *rctt*-**2d**/**1d** molar ratio is equal to 4.3 : 1 (degree of back transformation equals to 10%). Thus, we proved that both direct and back PCA reactions in the solid state of dyes **1** are feasible in principle.

### X-ray structural investigations

For all synthesized compounds, the X-ray structural study was performed. Single crystals of the dyes were grown by slow saturation of their acetonitrile solutions by benzene or benzene–hexane vapors in darkness. Another crystal modification of dye **1f** was prepared by slow evaporation of its solution in a water–MeCN mixture.

Structures of formula units for compounds **1c–g** are shown in Fig. 2. For dyes **1a** and **1b**, structural determination has been performed earlier.<sup>25,26</sup>

The independent moieties of crystal unit cells of **1a**, **b**, **d**, **e**, **g** and **1f**·0.5C<sub>6</sub>H<sub>6</sub> include one organic cation and one anion, whereas crystal unit cells of **1c**·0.5MeCN·1.5H<sub>2</sub>O and **1f** contain



Scheme 4 Synthesis of dyes **1c–g**.

Table 1 The starting dye **1**-to-cyclobutane derivative *rctt*-**2** ratio and the degree of conversion ( $\chi$ ) in polycrystalline films of dyes **1a–g** irradiated with visible light

Compound	<i>T</i> /h <sup>a</sup>	Dye : cyclobutane <sup>b</sup> (mol/mol)	$\chi$ <sup>b</sup> (%)
<b>1a</b> <sup>c</sup>	10	1 : 5.8	92
<b>1b</b> <sup>c</sup>	10	5.4 : 1	27
<b>1c</b>	10	1 : 18.5	97
<b>1d</b>	10	0 : 1	100
<b>1e</b>	10	1 : 3.3	87
	26	1 : 10.4	95
<b>1f</b>	10	2.77 : 1	42
	50	1.13 : 1	64
	240	1 : 1.34	73
<b>1g</b>	10	1 : 0	0

<sup>a</sup> The duration of irradiation. <sup>b</sup> Based on the <sup>1</sup>H NMR spectroscopic data. <sup>c</sup> Data from ref. 25.




Fig. 1  $^1\text{H}$  NMR spectra (aromatic protons region,  $\text{DMSO}-d_6$ ) of (a) dye **1d**, (b) dye **1d** irradiated with visible light in thin film (cyclobutane *rctt*-**2d**), and (c) *rctt*-**2d** irradiated with UV light in thin film (a 4.3:1 mixture of **2d** and **1d**; the proton signals of minor dye **1d** are marked with an asterisk).

two independent cations and two independent anions. The independent crystallographic unit for **1c** also contains one acetonitrile solvate molecule and three water molecules, one of which is disordered over two positions close to each other with nearly equal occupancies. The asymmetric unit of **1f**· $0.5\text{C}_6\text{H}_6$  contains half a benzene solvate molecule situated at a symmetry centre.

The cation in structure **1e** reveals a “pedal” disorder. This disorder is a consequence of a temperature-dependent dynamic process of “pedal” isomerisation in the crystal<sup>18,47–49</sup> when the ethylene fragment is subjected to a rotation about its single bonds (Scheme 6). Rather often this phenomenon is observed in styryl dyes<sup>25–27,29</sup> and in neutral styrylheterocycles.<sup>45</sup> The ratio of “pedal” isomers in **1e** corresponds to 0.64:0.36.

In crystal **1e** the  $\text{PF}_6^-$  anion is also disordered over two close positions. The rotational disorder of the  $\text{BF}_4^-$  anion over at least three positions is also observed for **1d**, with the ratio of occupancies 0.51:0.33:0.16. In reality, many residual peaks of the electron density are observed in the vicinity of the boron atom. However, their inclusion as partial atoms shows no improvement in the least-square refinement of the structure.

### Crystal packings and feasibility of PCA transformation

An analysis of the CSD data<sup>50</sup> made it possible to select six canonical packing forms for planar molecules (organic cations) of unsaturated compounds (Scheme 7).

Packings *a* and *b* are parallel-stacking and parquet-stacking, packings *c* and *d* are staircase and parquet, and packings *e* and *f* are dimeric-parallel and dimeric-parquet. In packings *c* and *d*,

there is no overlapping of the conjugated fragments. These packings can be deduced from packings *a* and *b*, respectively, by an equal slide to the same side of any moiety with respect to the previous one. In packings *e* and *f*, any dimeric  $\pi$ -conjugated moiety does not overlap in the projection with neighbouring ones. These packings may also be deduced from packings *a* and *b*, respectively, by an analogous slide of pairs of molecules.

These types of crystal packings are typical for styryl dyes, protonated styrylheterocycles  $\text{R-Het}^+-\text{CH}=\text{CH}-\text{Ar X}^-$  (Het – nitrogen-containing heterocycle,  $\text{X}^-$  – anion), and neutral styrylheterocycles  $\text{Het}-\text{CH}=\text{CH}-\text{Ar}$ . Packings *a*, *b*, *e*, and *f* may contain dimers preorganized for PCA reaction, whereas, packings *c* and *d* cannot contain such dimers. Our previous work showed that styryl dyes and protonated styrylheterocycles almost exclusively form crystal packings of *a* and *b* types, whereas, packings *c*, *d*, *f*, and *e* are typical for neutral styrylheterocycles.<sup>23–27,29</sup>

For ethylene derivatives prone to participate in PCA transformation, stacking packing motifs may be of different types depending on the symmetry operation combining structural units in a stack. Centrosymmetrically related stacks with the *syn*–“head-to-tail” organization of structural units form most frequently, whereas, translation-related stacks of the *syn*–“head-to-head” structure unit organization occur rather rarely.<sup>25–27,51,52</sup> Only one case of 2-fold-axis-related stacks is known for styryl dyes.<sup>26</sup>

In the case of translation-related stacks (Scheme 8), the PCA reaction in the crystal without its decomposition is impossible.


In this stack, one event of PCA would result in the loss of its local symmetry and defect formation. If the number of the events increases, the defect region rises. This eventually results in crystal degradation. In reality, the formation of this type of stacks in styryl dyes is associated with the significant slide of adjacent cations with respect to one another in order to reduce the Coulomb interaction between similarly charged fragments of adjacent cations. Therefore, the distance between the ethylene fragments of adjacent cations,  $d$ , is much longer than 4.2 Å and this also makes PCA transformation impossible in crystals of this type of packing.

In centrosymmetrically related stacks of the *syn*–“head-to-tail” type, the mutual organization of structural units is favorable for PCA. In this case, the PCA reaction is feasible, sometimes without single crystal degradation. A set of additional conditions is required for this.<sup>23–27,29</sup> The relationships  $d_1 < 4.2$  Å and  $d_2 > 4.2$  Å, where  $d_1$  and  $d_2$  are alternating distances between adjacent ethylene fragments in a stack, are often beneficial to PCA (Scheme 9).

In the case when both distances satisfy the condition of the PCA feasibility ( $d_1$  and  $d_2 < 4.2$  Å), the PCA process should go statistically between crystallographically independent pairs of structural units related by symmetry centres belonging to different crystallographic systems. This inevitably results in a general crystal symmetry violation, that is, crystal degradation.

The second additional condition is the occurrence of a soft flexible shell about the dimers, capable of reducing







**Fig. 2** Independent molecular moieties of crystal structures **1c–g**. Displacement ellipsoids are drawn at the 50% probability level.

cations or molecules, such as crown-ether fragments or bulky alkyl substituents.


In the crystal packing of  $1f \cdot 0.5C_6H_6$ , isolated alternating layers formed by stacks of cations and anions occur (Fig. 3).




**Scheme 6** The mechanism of the “pedal” motion in crystals of ethylene compounds.




**Scheme 7** Canonical packing motifs for planar structural units, molecules or cations; solid lines denote planar structure units in profile.



**Scheme 8** The translation-related stack of the *syn*-“head-to-head” type; *T* – translation, *d* – distance between adjacent ethylene fragments.




**Scheme 9** The centrosymmetrically related stack of the *syn*-“head-to-tail” type; *T* – translation, • – symmetry centre, *d*<sub>1</sub> and *d*<sub>2</sub> – distances between ethylene fragments in dimers and between the dimers, respectively.



**Fig. 3** The crystal packing of **1f**·0.5C<sub>6</sub>H<sub>6</sub>.

The layers of anions also involve benzene solvate molecules. The inclusion of large anions that are prone to form CTCs does not prevent the formation of stacks of the cations.

The mutual arrangement of cations in a stack (Fig. 4) is unfavorable for the PCA reaction to be accomplished in this crystal. Actually, ethylene fragments of the adjacent cations are not parallel. Moreover, they do not overlap in the projection (Fig. 4, (b)). The first of these unfavorable circumstances might be overcome due to the aforementioned “pedal” motion, and the second, because of mutual translation of molecules. Examples of a PCA reaction in the solid state



**Fig. 4** The structure of a cation stack in **1f**·0.5C<sub>6</sub>H<sub>6</sub> in two projections.

proceeding in spite of long distances ( $>5$  Å) between ethylene compounds were described earlier.<sup>47,53</sup> However, crystal **1f**·0.5C<sub>6</sub>H<sub>6</sub> manifests high stability to irradiation with visible light.

In the layer formed by picrate anions and benzene solvate molecules (Fig. 3), there are no  $\pi$ -stacking interactions between anions since they do not overlap in the projection. This layer does not contain conformationally flexible elements. Apparently, the structural rigidity of this layer explains high stability of crystal **1f**·0.5C<sub>6</sub>H<sub>6</sub> to irradiation.

The second crystal form of **1f** does not contain solvent molecules. Its crystal packing is different. It is composed of mixed stacks that involve both cations and anions. In a stack, four cations alternate with four anions (Fig. 5). Two central cations form a centrosymmetrically related couple with the *syn*-“head-to-tail” arrangement and the C6...C7A/C6A...C7 intermolecular distance equals to 3.79 Å, which is favorable for the PCA reaction that should result in cyclobutane *rcctt*-2f. Two other cations in the stack present crystallographically independent species. They are arranged in a pseudo-centrosymmetric mode with respect to the cations that form the central centrosymmetric couple. The intermolecular distance C6...C7' and C6'...C7 in the adjacent couple of cations (both 6.09 Å) is too long, which prevents the PCA transformation between these cations.




Fig. 5 A stack of cations and anions in crystal **1f**.

In the quaternion of anions, only two central ones are involved in a rather strong  $\pi$ -stacking interaction. The neighboring cation and anion overlap in the projection thus form CTC. This packing appears to predominate in a polycrystalline film of **1f** because it allows the PCA transformation that actually takes place, although relatively slowly (see Table 1).

Crystal **1c**·0.5MeCN·1.5H<sub>2</sub>O is built of alternating layers of cations and anions. The anion layers also involve water and acetonitrile solvate molecules. The sulfonate groups of the tosylate anions and water molecules are involved in a network of hydrogen bonds (see Fig. 2).

Stacks of cations are arranged in the *syn*-“head-to-head” pattern (Fig. 6). According to the aforementioned consideration, this type of stack organization excludes the feasibility of PCA in a single crystal. Actually, the distances between neighboring ethylene fragments are in the range of 7.62–8.22 Å that exceeds greatly the longest distance when the PCA reaction can proceed. Indeed, crystal **1c**·0.5MeCN·1.5H<sub>2</sub>O is stable to irradiation with visible light. However the PCA reaction occurs in the polycrystalline film of dye **1c** very efficiently to give the centrosymmetric isomer of cyclobutane derivative *rcctt*-2c. This means that the predominant packing motif in fine-grained crystals of the polycrystalline film of **1c** differs from that found in single crystals.

In the crystal packing of dye **1b**,<sup>25,26</sup> cations are arranged in stacks of the *syn*-“head-to-tail” type (Fig. 7). Any two adjacent cations are related *via* a symmetry centre. The distance between the ethylene fragments of cations A and D (C6A...C7D and C7A...C6D) is equal to 4.18 Å and lies in the vicinity of the upper limit of the PCA reaction feasibility.




Fig. 6 The stack of cations in structure **1c**·0.5MeCN·1.5H<sub>2</sub>O in two projections.




Fig. 7 The mutual arrangement of cations of **1b** in a stack.

While the distance between the ethylene fragments of the adjacent cation pair, D and B ( $C6D \cdots C7B$  and  $C7D \cdots C6B$ ) is too long, 6.89 Å, for the PCA reaction to proceed.

The mutual arrangement of cation pair A and D is not the best one for the feasibility of the PCA reaction (Fig. 8). The region of the  $\pi$ -overlap of the two cations covers only the periphery of their conjugated systems, which is not optimal for the PCA reaction in the single crystal. Apparently, this is defined in most degree by the influence of the  $I^-$  anion on the crystal packing. This anion tends to form secondary bonds of the  $C-H \cdots I^-$  type (Fig. 9), which results in shifts of the cations from the positions corresponding to the optimal geometry of preorganized dimers with an efficient  $\pi \cdots \pi$  overlap between the cations of the A and D type. Distances  $I1 \cdots H1$ ,  $I1 \cdots H4$ ,  $I1 \cdots H5$  are equal to 2.90, 3.24, 3.10 Å, which are shorter or equal to the minimum value of the van der Waals distance ( $\sim 3.2$  Å) and correspond to usual distances for the interactions of this type.<sup>54,55</sup> Involvement of cations in these weak interactions apparently deteriorates preorganization of the A $\cdots$ D type cations for PCA.

However, taking into account a high trend of ethylene groups to parallel displacement in the course of PCA, one may expect to accomplish this reaction in crystals of **1b**, but only with crystal degradation. Indeed, the PCA process is very slowly developing in crystals of **1b**. For two weeks, these crystals are stable to irradiation with visible light, but then




Fig. 8 The projection of a cation onto the plane of another cation in **1b**.




Fig. 9 Environment of the  $I^-$  anion in crystal **1b**.

they begin to slowly decompose. In a polycrystalline film this compound behaves analogously.

In crystals **1a**, **1d**, and **1e** with rotationally flexible anions ( $ClO_4^-$ ,  $BF_4^-$ , and  $PF_6^-$ , respectively), crystal packings qualitatively similar to that observed in **1b** are realized. All these packings are described by motif *b* in Scheme 7. As an example, the packing of **1a** is shown in Fig. 10. It is composed of centrosymmetric stacks with the *syn*-“head-to-tail” mutual organization of cations. This organization involves dimers favorable for PCA.

Although differences in crystal packings of **1a**, **1d**, and **1e** are quantitative, they determine the different behaviors of the compounds with respect to PCA in single crystals.

In the crystal packing of **1a**, both distances  $d_1$  and  $d_2$  (3.74 and 3.66 Å) satisfy conditions  $d_1$  and  $d_2 < 4.2$  Å. Therefore, PCA reaction is possible in this case, but with crystal decomposition. Indeed, upon irradiation with visible light, crystals decompose to an amorphous powder and the compound becomes lighter in colour. The result of this reaction is the formation of the centrosymmetric *rctt* isomer of cyclobutane derivative **2a**, from both crystallographically independent pairs of cations.

In crystal **1e**, analogous *syn*-“head-to-tail” stacks of cations occur, in which different “pedal” isomers are statistically distributed. Therefore, pairs of adjacent cations with the major-major, minor-minor, and major-minor mutual arrangement may exist simultaneously. In the first two cases, the adjacent ethylene fragments are antiparallel, which is favourable for PCA. In the third case, the mutual arrangement of these fragments is near perpendicular, which is unfavourable for PCA. However, the aforementioned dynamic “pedal” process may change the mutual orientation of the ethylene fragments in the third type pair to make the photoreaction possible.

Earlier,<sup>29</sup> we have established that the dynamic process of “pedal” isomerization in the crystal of protonated styrylpyridine **3** actually takes place. This crystal contains two crystallographically independent cations that are situated in parallel one above another. One of these cations reveals a “pedal” disorder (Scheme 10). Only the ethylene bond of the minor component of this disordered system is parallel to the ethylene bond of another ordered cation.





Fig. 10 The crystal packing of **1a** in two projections.







**Scheme 10** The structure formula of compound **3** and its dimer formed by two independent cations in the crystal.

Nevertheless, the full PCA transformation without crystal decomposition has been observed in the crystal, which is only possible if the “pedal” motion actually takes place in the crystal.

Fig. 11 and 12 show stacks with participation of cations of the major–major and minor–minor types in **1e**.



**Fig. 11** Stacks formed by major “pedal” isomers (top) and mutual projection of adjacent pairs of cations (bottom) in **1e**.



**Fig. 12** Stacks formed by minor “pedal” isomers (top) and mutual projection of adjacent pairs of cations (bottom) in **1e**.

In both cases, distances  $d_1$  and  $d_2$  are shorter than 4.2 Å, which gave evidence that the PCA transformation in this crystal is only possible with crystal decomposition. Indeed, crystals **1e** decompose under irradiation with visible light to give an amorphous powder. Under the same conditions, a polycrystalline film of **1e** almost completely transforms into cyclobutane derivative *rctt-2e*.


Kept under visible light, crystal **1d** becomes colourless very quickly (Fig. 13). Big crystals crack into smaller ones, but small crystals are mechanically stable; their shape stays unchanged.

These results indicate a PCA transformation without single crystal degradation in **1d** (Scheme 11).


The change in colour turned out to be distinct after approximately 20 minutes – the time required for the manipulation of crystals under a microscope before X-ray diffraction experiment starts. Because of a high rate of the PCA transformation in a single crystal, we failed to perform a very precise experiment for single crystal **1d**. The rotational disorder of the  $\text{BF}_4^-$  anion also reduces the accuracy of this X-ray study.

In crystal **1d** the cations are arranged in centrosymmetrical stacks of the *syn*–“head-to-tail” type (Fig. 14). Alternating distances  $d_1$  and  $d_2$  are equal to 3.67 and 3.97 Å.

Although both distances are shorter than 4.2 Å, the A–D pair is better preorganized for PCA than two crystallographically equivalent pairs B–D and A–C. It is seen from Fig. 14 that the ethylene groups at the central pair (A–D) are bent



**Fig. 13** Crystals of **1d** in perfluorinated oil on a glass support with a 2D millimeter scale: a and c before irradiation; b and d after irradiation with unfiltered light from a Fiber Optical Light Source (200 W, Meiji Techno) for 10 min (d) and 2 h (b). Photographs are taken using a EMZ-13RT microscope (Meiji Techno).



**Scheme 11** Direct and back PCA reactions in single crystal **1d**.




Fig. 14 A stack of cations in crystal 1d.

towards one another. Moreover, in pair A–D there are better geometric conditions for conjugated fragments efficient  $\pi$ -overlapping (Fig. 15). This gave basis to the suggestion that in crystal 1d, the PCA reaction will proceed only in crystallographically equivalent pairs. Therefore this reaction should not be accompanied by crystal decomposition.

Indeed, after 3 h irradiation with visible light of the same crystal of 1d that had been studied using X-ray diffraction analysis it turned colourless. The X-ray study established that the complete PCA transformation of dye 1d into cyclobutane derivative *rcctt*-2d had taken place without crystal decomposition (Fig. 16). The  $\text{BF}_4^-$  anion remains rotationally disordered over three positions.

It should be noted that in contrast to crystal 1d, crystal 2d turned out to be of rather good quality.




Fig. 15 The mutual projection of crystallographically nonequivalent pairs of cations in 1d.




Fig. 16 The structure of cyclobutane derivative *rcctt*-2d formed on irradiation of 1d.

A superposition of a pair of structure units of the initial compound 1d and corresponding structural units of the product 2d is shown in Fig. 17. It is seen that the cyclobutane moiety of 2d lies inside the pair of the initial cations. The shift of the boron atoms in 2d from their positions in 1d is equal to 0.21 Å. The  $\text{BF}_4^-$  anions in both compounds are slightly rotated with respect to one another and the ratio of their occupancies is slightly changed (0.51:0.33:0.16 in 1d, 0.60:0.20:0.20 in 2d). One may conclude that the changes are due to tuning of anions to a slightly changed crystal environment in the course of the PCA reaction.

Apparently these results explain why the PCA reaction in 1d proceeds so quickly and without crystal decomposition, at least in small crystals. For this reason, we tried to discover a photochemical reaction of *retro*-PCA, 2d  $\rightarrow$  1d.

The same single crystal of compound 2d was then subjected to unfiltered UV-vis radiation using a L8252 mercury-xenon lamp (Hamamatsu) at room temperature. Its spectrum contains a set of wavelengths. Among these are the ones initiating both the direct and the back photo-transformations. After 4 h irradiation of single crystal 2d, it has turned bright yellow again (crystal 4d). It should be noted that when UV-vis irradiation of the crystal was stopped, it became colourless in approximately 30 minutes of keeping on light at room temperature, but its colour remained unchanged at low temperature ( $\sim 150$  K) for several hours.




Fig. 17 Superposition of structure units of initial compound 1d (dash lines) and product 2d (solid lines); the fluorine atoms of anions in 1d and 2d are shown with green and violet colours, respectively.

In crystal **4d** we expected to observe the coexistence of both compounds, **1d** and **2d** (mixed **1d/2d** system), but with an *a priori* unknown ratio of their contents. Similar mixed systems were observed earlier, many times as the results of incomplete direct PCA reactions (for examples, see ref. 4,24,26,27,29). Nevertheless this result could be important in this particular case because it might indicate that the back photochemical reaction is also feasible for the single crystal in parallel with the direct PCA reaction. Note that in the polycrystalline film of *rcctt-2d*, the *retro*-PCA does take place, resulting in the appearance of 10% initial dye **1d** (see Fig. 1 and its discussion).

Meanwhile, the actual result of the irradiation of single crystal **2d** turned out to be unexpected (Fig. 18). Instead of mixed system **1d/2d**, the disordered cyclobutane system **4d** has appeared, with differently oriented cyclobutane. The ratio of two components of cyclobutane is 0.89:0.11 and both are centrosymmetric *rcctt* isomers. This result might be easily interpreted if molecules of dye **1d** formed *in situ* during irradiation contain the “pedal” disorder. In this case, structure **4d** will directly form from the intermediate mixed **1d/2d** system. As the “pedal” disorder is lacking in initial crystal **1d**, it is obvious that such a transformation of one cyclobutane moiety into another one requires the back transformation of **2d** into the initial compound together with its “pedal” isomerization. New-formed **1d** compound quickly transforms into the cyclobutane derivative again, being in a photostationary equilibrium. However, during its existence, it also transforms into another “pedal” isomer. On light, these two “pedal” isomers further convert into two kinds of cyclobutanes to give observed disordered crystal **4d**. Although the “pedal” disorder in initial crystals **1d** is lacking, it could appear in the **1d-2d** mixed crystal as a result of small displacements in the crystal packing due to photochemical transformations.

It should be noted that, in structures **1d**, **2d**, and **4d**, the  $\text{BF}_4^-$  anion is found to be rotationally disordered over three positions with the ratios of occupancies 0.51:0.33:0.16 in **1d**, 0.60:0.20:0.20 in **2d**, and 0.38:0.38:0.24 in **4d**. The ratios in **1d** and **2d** are rather close, whereas in **4d**, they are




Fig. 18 The structure of cyclobutane derivative **4d** obtained after 4 h UV-vis irradiation of crystal **2d**.

significantly different. This fact shows that the nearest environment of the reacting components of UV-irradiated crystal **2d** may differ enough from that in crystals **1d** and **2d** for making feasible the “pedal” motion in the crystal.

Thus, obtained structure **4d** serves as a proof that both photochemical processes – PCA and *retro*-PCA – are feasible in the same crystal without its decomposition.

In the crystal of dye **1g** containing a huge  $\text{BPh}_4^-$  anion, the main structure-forming role belongs to anions, which interrupt the cation's trend to form  $\pi$ -stacking elements in the crystal packing. That is, the main packing motifs for planar molecules shown in Scheme 7 are not applied here. This is in good agreement with the absence of PCA products in irradiated polycrystalline films of dye **1g** (see Table 1).

## Conclusions

This investigation made it possible to establish that the nature of the anion in styryl dyes affects significantly the crystal packing motif. Rotationally flexible anions such as  $\text{ClO}_4^-$ ,  $\text{BF}_4^-$ ,  $\text{PF}_6^-$  present the most favorable type for the realization of the centrosymmetrically related stacking motif of the organic cations. In most of the structures investigated, a trend to form stacking elements of the packing motif for cations is observed. Even the ability of picrate anion to form charge-transfer complexes with aromatic moieties does not violate this trend. Only the bulky  $\text{BPh}_4^-$  anion representing the major structure-forming element in the crystal of the corresponding dye is capable of suppressing this trend.

We have received the first proof that direct and back reactions of  $[2 + 2]$  photocycloaddition can be accomplished in the same single crystal of an ethylene compound with the retention of the single crystal. We believe that this process could be accomplished repeatedly in a crystal, which is important in creating switching systems or memory cells.

## Experimental section

### General

Melting points (uncorrected) were measured in capillaries using a Mel-Temp II apparatus.  $^1\text{H}$  NMR spectra were recorded using a Bruker DRX500 instrument (500.13 MHz) in  $\text{DMSO-d}_6$  at 25–30 °C using the solvent as an internal standard ( $\delta_{\text{H}}$  2.50 ppm). Elemental analyses were performed at the Laboratory of Microanalysis of A. N. Nesmeyanov Institute of Elementoorganic Compounds of the Russian Academy of Sciences (Moscow, Russian Federation). The samples for elemental analyses were dried at 80 °C *in vacuo*.

### Preparations

Syntheses of 4-[(*E*)-2-(3,4-dimethoxyphenyl)-1-ethenyl]pyridine, dyes **1a**, **b** and cyclobutanes *rcctt-2a*, **b** were described earlier.<sup>25,45</sup>

4-[(*E*)-2-(3,4-Dimethoxyphenyl)-1-ethenyl]-1-ethylpyridinium tosylate (**1c**). A mixture of 4-[(*E*)-2-(3,4-dimethoxyphenyl)-1-ethenyl]pyridine (0.31 g, 1.27 mmol) and ethyl 4-methylbenzenesulfonate (0.76 g, 3.81 mmol) was heated at



120 °C (oil bath) in darkness for 2 h. After cooling to room temperature, the resulting mass was washed with benzene (2 × 10 mL) and the solids were twice extracted with hot benzene (15 mL, for 1 h) in darkness. The insoluble substance was dissolved in abs. EtOH (3 mL) and the solution was diluted with Et<sub>2</sub>O (30 mL). The glassy precipitate thus formed was decanted and dried at 70 °C *in vacuo* in darkness to give dye **1c** (0.49 g, yield 87%) as a yellow-orange powder, mp 146–148 °C. Calcd. for C<sub>24</sub>H<sub>27</sub>NO<sub>5</sub>S·0.25H<sub>2</sub>O: C 64.63, H 6.21, N 3.14. Found: C 64.65, H 6.64, N 3.05%. <sup>1</sup>H NMR: δ = 1.53 (t, *J* = 7.3 Hz, 3H, MeCH<sub>2</sub>), 2.29 (s, 3H, MeAr), 3.83 (s, 3H, 4'-MeO), 3.85 (s, 3H, 3'-MeO), 4.51 (q, *J* = 7.3 Hz, 2H, CH<sub>2</sub>N), 7.08 (d, *J* = 8.4 Hz, 1H, 5'-H), 7.10 (d, *J* = 8.0 Hz, 2H, 3''-H, 5''-H), 7.30 (dd, *J* = 8.4 Hz, *J* = 1.5 Hz, 1H, 6'-H), 7.39 (br.s, 1H, 2'-H), 7.40 (d, *J* = 16.3 Hz, 1H, CH=CHPy), 7.48 (d, *J* = 8.0 Hz, 2H, 2''-H, 6''-H), 7.96 (d, *J* = 16.3 Hz, 1H, CH=CHPy), 8.16 (d, *J* = 6.7 Hz, 2H, 3-H, 5-H), 8.90 (d, *J* = 6.7 Hz, 2H, 2-H, 6-H).

**4-[(*E*)-2-(3,4-Dimethoxyphenyl)-1-ethenyl]-1-ethylpyridinium tetrafluoroborate (1d).** A solution of NaBF<sub>4</sub> (20 mg, 0.17 mmol) in MeOH (1 mL) was added to a solution of dye **1c** (64 mg, 0.15 mmol) in MeOH (1 mL) in darkness and the resulting mixture was cooled to 5 °C. The precipitate thus formed was filtered, washed with cold MeOH (2 × 1 mL) and dried in air to give dye **1d** (36 mg, yield 73%) as a yellow powder, mp 219–220 °C. Calcd. for C<sub>17</sub>H<sub>20</sub>BF<sub>4</sub>NO<sub>2</sub>: C 57.17, H 5.64, N 3.92. Found: C 57.24, H 5.67, N 3.88%. <sup>1</sup>H NMR: δ = 1.52 (t, *J* = 7.3 Hz, 3H, MeCH<sub>2</sub>), 3.83 (s, 3H, 4'-MeO), 3.85 (s, 3H, 3'-MeO), 4.50 (q, *J* = 7.3 Hz, 2H, CH<sub>2</sub>N), 7.08 (d, *J* = 8.3 Hz, 1H, 5'-H), 7.29 (dd, *J* = 8.3 Hz, *J* = 1.8 Hz, 1H, 6'-H), 7.39 (d, *J* = 1.8 Hz, 1H, 2'-H), 7.41 (d, *J* = 16.3 Hz, 1H, CH=CHPy), 7.97 (d, *J* = 16.3 Hz, 1H, CH=CHPy), 8.16 (d, *J* = 6.9 Hz, 2H, 3-H, 5-H), 8.91 (d, *J* = 6.9 Hz, 2H, 2-H, 6-H).

**4-[(*E*)-2-(3,4-Dimethoxyphenyl)-1-ethenyl]-1-ethylpyridinium hexafluorophosphate (1e).** A solution of NH<sub>4</sub>PF<sub>6</sub> (45 mg, 0.28 mmol) in MeOH (1 mL) was added to a solution of dye **1c** (102 mg, 0.23 mmol) in MeOH (2 mL) and the resulting mixture was cooled to 5 °C in darkness. The precipitate thus formed was filtered, washed with cold MeOH (2 × 1 mL) and dried in air to give dye **1e** (80 mg, yield 84%) as a yellow powder, mp 242–244 °C. Calcd. for C<sub>17</sub>H<sub>20</sub>F<sub>6</sub>NO<sub>2</sub>P: C 49.16, H 4.85, N 3.37. Found: C 49.06, H 4.84, N 3.29%. <sup>1</sup>H NMR: δ = 1.53 (t, *J* = 7.3 Hz, 3H, MeCH<sub>2</sub>), 3.83 (s, 3H, 4'-MeO), 3.85 (s, 3H, 3'-MeO), 4.51 (q, *J* = 7.3 Hz, 2H, CH<sub>2</sub>N), 7.08 (d, *J* = 8.3 Hz, 1H, 5'-H), 7.30 (dd, *J* = 8.3 Hz, *J* = 1.6 Hz, 1H, 6'-H), 7.38 (br.s, 1H, 2'-H), 7.40 (d, *J* = 16.3 Hz, 1H, CH=CHPy), 7.96 (d, *J* = 16.3 Hz, 1H, CH=CHPy), 8.16 (d, *J* = 6.7 Hz, 2H, 3-H, 5-H), 8.90 (d, *J* = 6.7 Hz, 2H, 2-H, 6-H).

**4-[(*E*)-2-(3,4-Dimethoxyphenyl)-1-ethenyl]-1-ethylpyridinium picrate (1f).** A solution of sodium picrate (42 mg, 0.17 mmol) in MeOH (1.5 mL) was added to a solution of dye **1c** (67 mg, 0.15 mmol) in MeOH (1 mL) and the resulting mixture was cooled to 5 °C in darkness. The precipitate thus formed was filtered, washed with cold MeOH (2 × 1 mL) and dried in air to give dye **1f** (57 mg, yield 75%) as a yellow powder, mp 149–150 °C. Calcd. for C<sub>23</sub>H<sub>22</sub>N<sub>4</sub>O<sub>9</sub>: C 55.42, H 4.45, N 11.24. Found: C 55.64, H 4.41, N 11.15%. <sup>1</sup>H NMR:

δ = 1.53 (t, *J* = 7.3 Hz, 3H, MeCH<sub>2</sub>), 3.83 (s, 3H, 4'-MeO), 3.85 (s, 3H, 3'-MeO), 4.51 (q, *J* = 7.3 Hz, 2H, CH<sub>2</sub>N), 7.08 (d, *J* = 8.4 Hz, 1H, 5'-H), 7.30 (dd, *J* = 8.4 Hz, *J* = 1.8 Hz, 1H, 6'-H), 7.38 (br.s, 1H, 2'-H), 7.40 (d, *J* = 16.3 Hz, 1H, CH=CHPy), 7.96 (d, *J* = 16.3 Hz, 1H, CH=CHPy), 8.16 (d, *J* = 6.8 Hz, 2H, 3-H, 5-H), 8.58 (s, 2H, 3''-H, 5''-H), 8.90 (d, *J* = 6.8 Hz, 2H, 2-H, 6-H).

**4-[(*E*)-2-(3,4-Dimethoxyphenyl)-1-ethenyl]-1-ethylpyridinium tetraphenylborate (1g).** A solution of sodium tetraphenylborate (146 mg, 0.43 mmol) in MeOH (2.5 mL) was added to a solution of dye **1c** (157 mg, 0.36 mmol) in MeOH (5 mL) and the resulting mixture was cooled to –10 °C in darkness. The precipitate thus formed was filtered, washed with cold MeOH (2 × 2 mL) and dried in air to give dye **1g** (206 mg, yield 98%) as a yellow powder, mp 195–197 °C. Calcd. for C<sub>41</sub>H<sub>40</sub>BNO<sub>2</sub>: C 83.53, H 6.84, N 2.38. Found: C 83.57, H 6.70, N 2.24%. <sup>1</sup>H NMR: δ = 1.52 (t, *J* = 7.3 Hz, 3H, MeCH<sub>2</sub>), 3.83 (s, 3H, 4'-MeO), 3.85 (s, 3H, 3'-MeO), 4.50 (q, *J* = 7.3 Hz, 2H, CH<sub>2</sub>N), 6.78 (t, *J* = 7.2 Hz, 4H, 4 4''-H), 6.92 (t, *J* = 7.4 Hz, 8H, 4 3''-H, 4 5''-H), 7.08 (d, *J* = 8.4 Hz, 1H, 5'-H), 7.17 (m, 8H, 4 2''-H, 4 6''-H), 7.29 (dd, *J* = 8.4 Hz, *J* = 1.8 Hz, 1H, 6'-H), 7.38 (d, *J* = 1.8 Hz, 1H, 2'-H), 7.40 (d, *J* = 16.3 Hz, 1H, CH=CHPy), 7.96 (d, *J* = 16.3 Hz, 1H, CH=CHPy), 8.16 (d, *J* = 6.8 Hz, 2H, 3-H, 5-H), 8.90 (d, *J* = 6.8 Hz, 2H, 2-H, 6-H).

#### Synthesis of cyclobutane derivatives from dyes **1** (general procedure)

A solution of dye **1c–g** (40 μmol) in MeCN (~0.5 mL) was concentrated in a 10 cm Petri dish to form a thin polycrystalline film of the corresponding compound. The sample was irradiated with unfiltered light using a 60 W incandescent lamp at a distance of 15 cm for 10–240 h. The resulting compounds were collected mechanically. The compositions of the products and the degree of conversion into cyclobutane derivatives were analyzed based on the <sup>1</sup>H NMR spectroscopy data (in DMSO-*d*<sub>6</sub>) by comparing the integrated intensities of the signals for protons. The experimental data obtained at different exposure times are given in Table 1. In the cases when the degree of conversion was ≥95%, the characteristics of the resulting cyclobutane derivatives are given.

**4,4'-[2-*c*,4-*t*-Bis(3,4-dimethoxyphenyl)cyclobutane-1-*r*,3-*t*-diyl]bis(1-ethylpyridinium) ditosylate (*rcctt*-2c).** Compound *rcctt*-2c (11.4 mg, yield 65%) was obtained as a yellowish powder, mp 70–75 °C. Calcd. for C<sub>48</sub>H<sub>54</sub>N<sub>2</sub>O<sub>10</sub>S<sub>2</sub>·1.5H<sub>2</sub>O: C 63.35, H 6.31, N 3.08. Found: C 63.49, H 6.27, N 3.13%. <sup>1</sup>H NMR: δ = 1.41 (t, *J* = 7.3 Hz, 6H, 2 MeCH<sub>2</sub>), 2.29 (s, 6H, 2 MeAr), 3.65 (s, 6H, 2 3'-MeO), 3.67 (s, 6H, 2 4'-MeO), 4.49 (q, *J* = 7.3 Hz, 4H, 2 CH<sub>2</sub>N), 4.81 (dd, *J* = 9.8 Hz, *J* = 7.3 Hz, 2H, 2 CHAr), 4.90 (dd, *J* = 9.8 Hz, *J* = 7.3 Hz, 2H, 2 CHPy), 6.77 (s, 4H, 2 5'-H, 2 6'-H), 6.80 (s, 2H, 2 2'-H), 7.11 (d, *J* = 8.0 Hz, 4H, 2 3''-H, 2 5''-H), 7.47 (d, *J* = 8.0 Hz, 4H, 2 2''-H, 2 6''-H), 7.95 (d, *J* = 6.7 Hz, 4H, 2 3-H, 2 5-H), 8.88 (d, *J* = 6.7 Hz, 4H, 2 2-H, 2 6-H).

**4,4'-[2-*c*,4-*t*-Bis(3,4-dimethoxyphenyl)cyclobutane-1-*r*,3-*t*-diyl]bis(1-ethylpyridinium) di(tetrafluoroborate) (*rcctt*-2d).** Compound *rcctt*-2d (11.6 mg, yield 85%) was obtained as a



slightly yellowish powder, mp 194–196 °C. Calcd. for  $C_{34}H_{40}B_2F_8N_2O_4$ : C 57.17, H 5.64, N 3.92. Found: C 57.31, H 5.69, N 3.98%.  $^1H$  NMR:  $\delta$  = 1.42 (t,  $J$  = 7.3 Hz, 6H, 2 MeCH<sub>2</sub>), 3.65 (s, 6H, 2 3'-MeO), 3.67 (s, 6H, 2 4'-MeO), 4.49 (q,  $J$  = 7.3 Hz, 4H, 2 CH<sub>2</sub>N), 4.81 (dd,  $J$  = 9.6 Hz,  $J$  = 7.4 Hz, 2H, 2 CHAr), 4.89 (dd,  $J$  = 9.6 Hz,  $J$  = 7.4 Hz, 2H, 2 CHPy), 6.77 (s, 4H, 2 5'-H, 2 6'-H), 6.80 (s, 2H, 2 2'-H), 7.95 (d,  $J$  = 6.7 Hz, 4H, 2 3-H, 2 5-H), 8.88 (d,  $J$  = 6.7 Hz, 4H, 2 2-H, 2 6-H).

**4,4'-[2-*c*,4-*t*-Bis(3,4-dimethoxyphenyl)cyclobutane-1-*r*,3-*t*-diyl]bis(1-ethylpyridinium) di(hexafluorophosphate) (*rc*tt-2e).** Compound *rc*tt-2e (15.1 mg, yield 90%) was obtained as a yellowish powder, mp 225–230 °C. Calcd. for  $C_{34}H_{40}F_{12}N_2O_4P_2$ : C 49.16, H 4.85, N 3.37. Found: C 49.09, H 4.86, N 3.35%.  $^1H$  NMR:  $\delta$  = 1.42 (t,  $J$  = 7.3 Hz, 6H, 2 MeCH<sub>2</sub>), 3.65 (s, 6H, 2 3'-MeO), 3.67 (s, 6H, 2 4'-MeO), 4.49 (q,  $J$  = 7.3 Hz, 4H, 2 CH<sub>2</sub>N), 4.81 (dd,  $J$  = 9.7 Hz,  $J$  = 8.3 Hz, 2H, 2 CHAr), 4.89 (dd,  $J$  = 9.7 Hz,  $J$  = 8.3 Hz, 2H, 2 CHPy), 6.77 (s, 4H, 2 5'-H, 2 6'-H), 6.80 (s, 2H, 2 2'-H), 7.95 (d,  $J$  = 6.7 Hz, 4H, 2 3-H, 2 5-H), 8.88 (d,  $J$  = 6.7 Hz, 4H, 2 2-H, 2 6-H).

### Photolysis of thin films of dye 1d and cyclobutane *rc*tt-2d

A solution of dye 1d (1.7 mg, 5  $\mu$ mol) in MeCN (0.1 mL) was concentrated in a Petri dish to form a thin polycrystalline film of this dye (spot diameter ~2 cm). The sample was irradiated

with the light from a L8253 xenon lamp (Hamamatsu, maximum power, ZhS10 light filter (transmission >390 nm), distance to the light source 10 cm) for 1 h. A half of the film was collected mechanically and then analyzed by  $^1H$  NMR method (in DMSO-*d*<sub>6</sub>), which showed full conversion into cyclobutane derivative *rc*tt-2d. The residual film consisting of *rc*tt-2d was irradiated for 2 h under the same conditions but using a set of ZhS3 and UFS5 light filters (transmission 280–340 nm). By comparing the integrated intensities of the signals for protons, analysis of the  $^1H$  NMR spectroscopy data of this film (as a solution in DMSO-*d*<sub>6</sub>) revealed 10% conversion into initial dye 1d. Fragments of these  $^1H$  NMR spectra are shown in Fig. 1.

### X-ray diffractometry

Single crystals of compounds 1c–g, 2d, and 4d were subjected to X-ray single crystal measurements using a Bruker CCD SMART diffractometer (SMART-6K for 1c–e, g and 1f–0.5C<sub>6</sub>H<sub>6</sub>, SMART-1K for 2d and SMART-APEX-II for 4d) with graphite monochromatized MoK $\alpha$  radiation (wavelength 0.71073 Å) and  $\omega$  scan mode, and a Bruker PLATINUM135 CCD area diffractometer for 1f with CuK $\alpha$  radiation (wavelength 1.54178 Å) and  $\phi$ – $\omega$  scan mode under a stream of cooled nitrogen gas. The reduction of experimental reflections was performed using the SAINT program.<sup>56</sup>

**Table 2** X-ray structure determination summary

Compound	1c·0.5MeCN·1.5H <sub>2</sub> O	1e	1f·0.5C <sub>6</sub> H <sub>6</sub>	1f	1g	1d	2d	4d
Empirical formula	C <sub>25</sub> H <sub>31.5</sub> N <sub>1.5</sub> O <sub>6.5</sub> S	C <sub>17</sub> H <sub>21</sub> F <sub>6</sub> N <sub>2</sub> O <sub>2</sub> P	C <sub>26</sub> H <sub>25</sub> N <sub>4</sub> O <sub>9</sub>	C <sub>23</sub> H <sub>22</sub> N <sub>4</sub> O <sub>9</sub>	C <sub>41</sub> H <sub>40</sub> BN <sub>2</sub> O <sub>2</sub>	C <sub>17</sub> H <sub>20</sub> BF <sub>4</sub> N <sub>2</sub> O <sub>2</sub>	C <sub>34</sub> H <sub>40</sub> B <sub>2</sub> F <sub>8</sub> N <sub>2</sub> O <sub>4</sub>	C <sub>34</sub> H <sub>40</sub> B <sub>2</sub> F <sub>8</sub> N <sub>2</sub> O <sub>4</sub>
<i>M</i>	489.08	415.31	537.50	498.45	589.55	357.15	714.30	714.30
Crystal system	Monoclinic	Monoclinic	Monoclinic	Monoclinic	Orthorhombic	Monoclinic	Monoclinic	Monoclinic
Space group	<i>P</i> 2 <sub>1</sub> / <i>n</i>	<i>P</i> 2 <sub>1</sub> / <i>c</i>	<i>P</i> 2 <sub>1</sub> / <i>c</i>	<i>P</i> 2 <sub>1</sub> / <i>c</i>	<i>P</i> 2 <sub>1</sub> 2 <sub>1</sub> 2 <sub>1</sub>	<i>P</i> 2 <sub>1</sub> / <i>c</i>	<i>P</i> 2 <sub>1</sub> / <i>c</i>	<i>P</i> 2 <sub>1</sub> / <i>c</i>
<i>a</i> /Å	15.6904(6)	7.106(2)	11.7370(10)	14.2243(10)	8.9861(4)	7.3589(3)	7.3008(3)	7.319(2)
<i>b</i> /Å	21.8223(8)	22.507(7)	26.623(2)	14.4190(9)	9.0343(4)	20.8300(9)	20.8807(6)	20.852(7)
<i>c</i> /Å	15.8024(6)	11.576(3)	8.2528(7)	22.4404(15)	40.1047(17)	11.2706(5)	11.3281(4)	11.302(4)
$\alpha$ /°	90	90	90	90	90	90	90	90
$\beta$ /°	113.2560(10)	90.143(8)	109.714(3)	104.933(4)	90	91.727(13)	91.018(2)	91.835(5)
$\gamma$ /°	90	90	90	90	90	90	90	90
<i>V</i> /Å <sup>3</sup>	4971.1(3)	1851.5(10)	2427.6(4)	4447.1(5)	3255.8(2)	1726.84(13)	1725.95(11)	1723.9(9)
<i>Z</i>	8	4	4	8	4	4	2	2
<i>F</i> (000)	2080	860	1124	2080	1256	744	744	744
$\mu$ /mm <sup>−1</sup>	0.174	0.219	0.113	0.991	0.072	0.117	0.117	0.117
<i>T</i> /K	120	120	120	100	120	120	120	150
Data collected	36 155	14 757	22 602	14 627	29 245	12 462	13 052	17 378
Unique data ( <i>R</i> <sub>int</sub> )	10 832 (0.1497)	4884 (0.0977)	5846 (0.1896)	4653 (0.0443)	9470 (0.0961)	4142 (0.0792)	4538 (0.0486)	4156 (0.0357)
$\theta$ range/°	1.55–27.00	0.90–28.99	1.84–26.5	3.22–53.37	2.03–30.00	1.96–28.00	1.95–29.00	2.05–28.00
No. of variables	634	409	356	649	409	237	318	302
<i>R</i> <sub>1</sub>	0.0712	0.0632	0.0808	0.0840	0.0732	0.0790	0.0572	0.0601
[ <i>I</i> > 2 $\sigma$ ( <i>I</i> )] <i>wR</i> <sub>2</sub>	0.1754	0.1431	0.1869	0.2049	0.1130	0.1786	0.1336	0.1668
(all data) GOF	0.868	1.168	0.914	1.046	0.962	1.042	1.045	1.093
$\Delta\rho_{\max}$ , min/e Å <sup>−3</sup>	0.629/−0.427	0.243/−0.302	0.464/−0.372	0.823/−0.363	0.405/−0.237	0.409/−0.323	0.489/−0.451	0.591/−0.372

The structures were solved by direct methods and refined on  $F^2$  by full-matrix least-squares in anisotropic approximation for non-hydrogen atoms using the SHELXTL-Plus<sup>57</sup> and Olex-2(ref. 58) software. Positions of hydrogen atoms were calculated geometrically. In experiments for **1c**, **1d**, **1e**, **1f**, **1f**·0.5C<sub>6</sub>H<sub>6</sub>, **1g** and **4d**, hydrogen atoms were refined using a “riding” model and for **2d** hydrogen atoms were refined in the isotropic approximation. The BF<sub>4</sub><sup>−</sup> anion in structures **1d**, **2d**, **4d** reveals a rotational disorder over three positions, with occupancies ratios discussed above. The PF<sub>6</sub><sup>−</sup> anion in **1e** is disordered over two positions, with one common fluorine atom and close positions of the P atoms. The ratio of occupancies of two components of the disorder is equal to 0.77:0.23. The cation in **1e** reveals the “pedal” disorder with the ratio of occupancies equal to 0.61:0.39.

For compound **1f** (unsolvated form), only very small single crystals visible in the microscope as bright points were obtained. Such single crystals gave only a few very weak X-ray reflections at Mo-radiation and rather weak ones at Cu-radiation. However, we have managed to collect a big number of rather weak experimental reflections in the near area of  $\theta$ . The structure was successfully solved and refined in the anisotropic approximation, but with rather high standard deviations in geometric parameters. However, taking into account that only the packing motif of this crystal is required for our discussion and also special importance of this particular structure to establish a correlation between its packing motif and the photochemical behaviour of the compound in thin polycrystalline films, we included these data in this article.

A summary of the crystallographic data and structure determination parameters is provided in Table 2.

The experimental data for all structures are deposited in the Cambridge Crystallographic Data Centre (CCDC registration numbers are 969548 (**1c**·0.5MeCN·1.5H<sub>2</sub>O), 859813 (**1d**), 969549 (**1e**), 969551 (**1f**), 969550 (**1f**·0.5C<sub>6</sub>H<sub>6</sub>), 969552 (**1g**), 859814 (**2d**), and 859815 (**4d**)).

## Acknowledgements

This work was supported by the Russian Foundation for Basic Research (project no. 14-03-00012), the Presidium of the Russian Academy of Sciences (program no. 9P), and the Photochemistry Centre of the Russian Academy of Sciences. L.G.K. and A.V.C. thank the Royal Society of Chemistry for International Authors Grants.

## Notes and references

- G. J. M. Schmidt, *Pure Appl. Chem.*, 1971, 27, 647.
- V. Ramamurthy and K. Venkatesan, *Chem. Rev.*, 1987, 87, 433.
- L. R. MacGillivray, *CrystEngComm*, 2002, 4(7), 37.
- I. Turowska-Tyrk, *J. Phys. Org. Chem.*, 2004, 17, 837.
- C.-H. Huang and D. M. Bassani, *Eur. J. Org. Chem.*, 2005, 4041.
- T. Friščić and L. R. MacGillivray, *Supramol. Chem.*, 2005, 17, 47.
- M. Nagarathinam and J. J. Vittal, *Macromol. Rapid Commun.*, 2006, 27, 1091.
- J. Svoboda and B. König, *Chem. Rev.*, 2006, 106, 5413.
- D.-K. Bučar, G. S. Papaefstathiou, T. D. Hamilton, Q. L. Chu, I. G. Georgiev and L. R. MacGillivray, *Eur. J. Inorg. Chem.*, 2007, 4559.
- S. P. Gromov, *Russ. Chem. Bull., Int. Ed.*, 2008, 57, 1325.
- N. Hoffmann, *Chem. Rev.*, 2008, 108, 1052.
- L. R. MacGillivray, *J. Org. Chem.*, 2008, 73, 3311.
- L. R. MacGillivray, G. S. Papaefstathiou, T. Friščić, T. D. Hamilton, D.-K. Bučar, Q. Chu, D. B. Varshney and I. G. Georgiev, *Acc. Chem. Res.*, 2008, 41, 280.
- M. H. Mir, L. L. Koh, G. K. Tan and J. J. Vittal, *Angew. Chem., Int. Ed.*, 2010, 49, 390.
- E. Elacqua and L. R. MacGillivray, *Eur. J. Org. Chem.*, 2010, 6883.
- B. R. Bhogala, B. Captain, A. Parthasarathy and V. Ramamurthy, *J. Am. Chem. Soc.*, 2010, 132, 13434.
- Y. Sonoda, *Molecules*, 2011, 16, 119.
- E. Elacqua, P. Kaushik, R. H. Groeneman, J. C. Sumrak, D.-K. Bučar and L. R. MacGillivray, *Angew. Chem., Int. Ed.*, 2012, 51, 1037.
- R. Santra, M. Garai, D. Mondal and K. Biradha, *Chem. – Eur. J.*, 2013, 19, 489.
- S. Bhattacharya, J. Stojaković, B. K. Saha and L. R. MacGillivray, *Org. Lett.*, 2013, 15, 744.
- F. Li, J. Zhuang, G. Jiang, H. Tang, A. Xia, L. Jiang, Y. Song, Y. Li and D. Zhu, *Chem. Mater.*, 2008, 20, 1194.
- A. Papagni, P. Del Buttero, C. Bertarelli, L. Miozzo, M. Moret, M. T. Pryce and S. Rizzato, *New J. Chem.*, 2010, 34, 2612.
- A. I. Vedernikov, S. P. Gromov, N. A. Lobova, L. G. Kuz'mina, Yu. A. Strelenko, J. A. K. Howard and M. V. Alfimov, *Russ. Chem. Bull., Int. Ed.*, 2005, 54, 1954.
- L. G. Kuz'mina, A. I. Vedernikov, N. A. Lobova, A. V. Churakov, J. A. K. Howard, M. V. Alfimov and S. P. Gromov, *New J. Chem.*, 2007, 31, 980.
- A. I. Vedernikov, L. G. Kuz'mina, S. K. Sazonov, N. A. Lobova, P. S. Loginov, A. V. Churakov, Yu. A. Strelenko, J. A. K. Howard, M. V. Alfimov and S. P. Gromov, *Russ. Chem. Bull., Int. Ed.*, 2007, 56, 1860.
- L. G. Kuz'mina, A. I. Vedernikov, S. K. Sazonov, N. A. Lobova, P. S. Loginov, J. A. K. Howard, M. V. Alfimov and S. P. Gromov, *Crystallogr. Rep.*, 2008, 53, 428.
- L. G. Kuz'mina, A. I. Vedernikov, J. A. K. Howard, M. V. Alfimov and S. P. Gromov, *Nanotechnologies in Russia*, 2008, 3, 408.
- S. P. Gromov, A. I. Vedernikov, L. G. Kuz'mina, D. V. Kondratuk, S. K. Sazonov, Yu. A. Strelenko, M. V. Alfimov and J. A. K. Howard, *Eur. J. Org. Chem.*, 2010, 2587.
- L. G. Kuz'mina, A. I. Vedernikov, S. K. Sazonov, N. A. Lobova, A. V. Churakov, E. Kh. Lermontova, J. A. K. Howard, M. V. Alfimov and S. P. Gromov, *Russ. Chem. Bull., Int. Ed.*, 2011, 60, 1734.

- 30 S. P. Gromov, A. I. Vedernikov, Yu. V. Fedorov, O. A. Fedorova, E. N. Andryukhina, N. E. Shepel', Yu. A. Strelenko, D. Johnels, U. Edlund, J. Saltiel and M. V. Alfimov, *Russ. Chem. Bull., Int. Ed.*, 2005, **54**, 1569.
- 31 M. V. Alfimov, S. P. Gromov, O. B. Stanislavskii, E. N. Ushakov and O. A. Fedorova, *Russ. Chem. Bull.*, 1993, **42**, 1385.
- 32 S. P. Gromov, O. A. Fedorova, E. N. Ushakov, A. V. Buevich and M. V. Alfimov, *Russ. Chem. Bull.*, 1995, **44**, 2131.
- 33 E. N. Ushakov, S. P. Gromov, A. V. Buevich, I. I. Baskin, O. A. Fedorova, A. I. Vedernikov, M. V. Alfimov, B. Eliasson and U. Edlund, *J. Chem. Soc., Perkin Trans. 2*, 1999, 601.
- 34 S. P. Gromov, N. A. Lobova, A. I. Vedernikov, L. G. Kuz'mina, J. A. K. Howard and M. V. Alfimov, *Russ. Chem. Bull., Int. Ed.*, 2009, **58**, 1211.
- 35 S. P. Gromov, A. I. Vedernikov, N. A. Lobova, L. G. Kuz'mina, S. S. Basok, Yu. A. Strelenko, M. V. Alfimov and J. A. K. Howard, *New J. Chem.*, 2011, **35**, 724.
- 36 H. Shayira Banu, A. Lalitha, K. Pitchumani and C. Srinivasan, *Chem. Commun.*, 1999, 607.
- 37 K. S. S. P. Rao, S. M. Hubig, J. N. Moorthy and J. K. Kochi, *J. Org. Chem.*, 1999, **64**, 8098.
- 38 S. Y. Jon, Y. H. Ko, S. H. Park, H.-J. Kim and K. Kim, *Chem. Commun.*, 2001, 1938.
- 39 D. G. Amirsakis, M. A. Garcia-Garibay, S. J. Rowan, J. F. Stoddart, A. J. P. White and D. J. Williams, *Angew. Chem., Int. Ed.*, 2001, **40**, 4256.
- 40 M. Pattabiraman, A. Natarajan, R. Kaliappan, J. T. Mague and V. Ramamurthy, *Chem. Commun.*, 2005, 4542.
- 41 R. Kaliappan, L. S. Kaanumalle, A. Natarajan and V. Ramamurthy, *Photochem. Photobiol. Sci.*, 2006, **5**, 925.
- 42 G. Wenz, B.-H. Han and A. Müller, *Chem. Rev.*, 2006, **106**, 782.
- 43 R. Kaliappan, M. V. S. N. Maddipatla, L. S. Kaanumalle and V. Ramamurthy, *Photochem. Photobiol. Sci.*, 2007, **6**, 737.
- 44 M. V. S. N. Maddipatla, L. S. Kaanumalle, A. Natarajan, M. Pattabiraman and V. Ramamurthy, *Langmuir*, 2007, **23**, 7545.
- 45 L. G. Kuz'mina, A. I. Vedernikov, N. A. Lobova, S. K. Sazonov, S. S. Basok, J. A. K. Howard and S. P. Gromov, *Russ. Chem. Bull., Int. Ed.*, 2009, **58**, 1192.
- 46 G. R. Desiraju, *J. Am. Chem. Soc.*, 2013, **135**, 9952.
- 47 G. S. Murthy, P. Arjunan, K. Venkatesan and V. Ramamurthy, *Tetrahedron*, 1987, **43**, 1225.
- 48 J. Harada and K. Ogawa, *J. Am. Chem. Soc.*, 2004, **126**, 3539.
- 49 J. Harada and K. Ogawa, *Chem. Soc. Rev.*, 2009, **38**, 2244.
- 50 F. H. Allen, *Acta Crystallogr., Sect. B: Struct. Sci.*, 2002, **58**, 380.
- 51 G. K. Kole, G. K. Tan and J. J. Vittal, *CrystEngComm*, 2012, **14**, 7438.
- 52 G. K. Kole, G. K. Tan and J. J. Vittal, *Org. Lett.*, 2010, **12**, 128.
- 53 *Photochemistry in Organized and Constrained Media*, ed. V. Ramamurthy, VCH, New York, 1991.
- 54 T. Steiner, *Acta Crystallogr., Sect. B: Struct. Sci.*, 1998, **54**, 456.
- 55 D. A. Dickie, D. Abeysekera, I. D. McKenzie, H. A. Jenkins and J. A. C. Clyburne, *Cryst. Eng.*, 2003, **6**, 79.
- 56 *SAINT, Version 6.02A*, Bruker AXS Inc., Madison, Wisconsin (USA), 2001.
- 57 *SHELXTL-Plus, Version 5.10*, Bruker AXS Inc., Madison, Wisconsin (USA), 1997.
- 58 O. V. Dolomanov, L. J. Bourhis, R. J. Gildea, J. A. K. Howard and H. Pushman, *J. Appl. Crystallogr.*, 2009, **42**, 339.